arXiv:0806.4947v2 [math.NT] 30 Jun 2008

Examples of finite p-divisible sets of MHS

Department of Mathematics, Eckerd College, St. Petersburg, FL 33711, USA

Jianqiang Zhao

The following examples give evidence to the following conjecture contained in my paper [2]. All the main theoretical results can be found in that paper.

Conjecture 1. *Let d be a positive integer and $\vec{s} \in \mathbb{N}^d$. Then the set $J(\vec{s}|p)$ is finite for every prime p.*

Example 2. The first example we would like to do is about the partial sums of $\zeta(2)$ series. The prime $p = 7$ is a little different from the others because $H(2;3) = 49/36$. We find that

$$J_1(2|7) = \{3,6\}, J_1(2|7^2) = \{3\}.$$

By [2, Thm 3.6] $n = 7\tilde{n} + r \in J_2(2|7)$, $0 \le r \le 6$, exists if and only if $\tilde{n} = 3$, and

$$H(r) \equiv -\psi(3) \equiv -1/36 \equiv -1 \pmod 7.$$

From the congruence

$$H(2;1) \equiv 1, \quad H(2;2) \equiv 3, \quad H(2;4) \equiv 4, \quad H(2;5) \equiv -1 \pmod 7,$$

we find $r = 5$. But it's an easy matter to check that

$$H(2;26) \equiv 14 \not\equiv 0 \pmod{7^2}.$$

Hence $J_2(2|7^2) = \emptyset$ and consequently $J(2|7) = \{0,3,6,26\}$.

For all the other primes $p \ne 7$ from 5 up to 1061[1] we find that

$$J_1(2|p) = \{0,(p-1)/2,p-1\} \cup T(2|p) \cup \{p-1-r : r \in T(2|p)\}$$

where $T(2|p)$ are listed in Table 1 if $T(2|p) \ne \emptyset$. Moreover, $J_1(2|p^2) = \emptyset$ which implies $J(2|p) = \{(p-1)/2, p-1\} \cup T(2|p) \cup \{p-1-r : r \in T(2|p)\}$ in this range.

p	$T(2\|p)$	p	$T(2\|p)$	p	$T(2\|p)$	p	$T(2\|p)$	p	$T(2\|p)$	p	$T(2\|p)$
37	15	163	61	419	111	563	175,227	677	153	883	151
41	4	167	61	421	59	569	199	709	123	911	345
43	11	181	85	433	179	571	247	727	197,239	929	64
59	6,24	211	99	457	216	577	134,158	739	93	953	199
97	15	241	60,96	467	158,170	601	17	787	344	967	463
107	39	269	50	479	5	617	97	797	185,226	971	429
127	23	307	27	487	32	619	16,70,286	811	371	991	72
137	44	311	43	491	173	643	17,222	821	39	997	205,310
149	37	373	54	499	134	653	246,307	859	414	1013	430
157	25	383	150	547	165	659	232	863	226	1031	384

Table 1: The set $J(2|p)$, $p \le 1061$, is finite.

---

[1] We choose 1061 because it's the first irregular prime greater than 1000.

Example 3. In this example we will determine $J(3|p)$ for primes p no greater than 1061. First let $p = 3$. Easy computation by hand will show that $J_1(3|3) = \{2\}$ and $J_1(3|3^3) = \emptyset$. Thus $J(3|3) = \{0, 2\}$. Let $p = 5$. We see that $J_1(3|5) = \{4\}$ and $J_1(3|5^3) = \emptyset$. Hence $J(3|5) = \{0, 4\}$.

Similarly, for p between 7 and 1061 we get that $J(3|p) = \{0, p-1\} \cup T(3|p) \cup \{p-1-r : r \in T(3|p)\}$ where $T(3|p)$ are listed in Table 2 if $T(3|p) \neq \emptyset$. In fact we find that when p lies in this range $J(3|p^2) = \{0, p-1\}$ exactly, and $J(3|p^3) = \{0\}$ except for $J(3|37^3) = \{0, 36\}$. But by our [1, Thm 2.6] we know that there might be infinitely many other primes with $J(3|p^3) = \{0, p-1\}$, as long as $(p, p-5)$ is an irregular pair.

Let's look at $p = 37$. Maple tells us that $J_1(3|37) = \{4, 13, 23, 32, 36\}$, $J_1(3|37^2) = J_1(3|37^3) = \{36\}$. By [2, Thm 3.6], in order for $n = 37\tilde{n} + r \in J_2(3|37)$ to exist it is necessary and sufficient that

$$\tilde{n} = 36 \text{ and } H(3, r) \equiv -\psi(\tilde{n}) \equiv 28 \pmod{37}.$$

A quick Maple computation picks up $r = 8$ and $r = 28$. (Recall that all such r should be paired and add up to $p - 1 = 36$). So $J_2(3|37) = \{37 \cdot 36 + 8, 37 \cdot 36 + 28\} = \{1340, 1360\}$. Direct computation shows that these two number do not lie in $J_2(3|37^3)$ which shows that $J_3(3|37) = \emptyset$. To summarize we get

$$J(3|37) = \{0, 4, 13, 23, 32, 36, 1340, 1360\}. \tag{1}$$

p	$T(3\mid p)$	p	$T(3\mid p)$	p	$T(3\mid p)$	p	$T(3\mid p)$	p	$T(3\mid p)$	p	$T(3\mid p)$
11	4	163	73,76	277	56	421	76	619	138	863	135
17	7	173	56	281	45	431	11,25	673	147	877	73
31	8	179	24	283	113	443	105	677	324	883	195
37	see (1)	181	76	317	103	479	127	743	307	887	145
47	5	193	38	331	154,161	521	43	751	52	911	287
53	6	223	7	347	16,73,107	523	93	773	311	941	241,419
67	28,30	227	91	349	83	563	62,156	787	9,199	953	29,69
89	10,43	251	3	359	10,136	571	274	809	213	1013	345
113	39,43	263	76	367	163	577	76	823	37	1021	383
137	44	269	131	379	160	599	224	829	256	1051	284
149	35	271	70,105	389	63	607	42	839	71	1061	515

Table 2: The set $J(3|p) = \{0, p-1\} \cup T(3|p) \cup \{p-1-r : r \in T(3|p)\}$, $p \leq 1061$, is finite.

Example 4. The partial sums of $\zeta(4)$ series are very regular. When $p = 3$ and $p = 5$, we see that

$$J_1(4|p) = \emptyset \Longrightarrow J(4|p) = \{0\}.$$

For all the primes from 7 up to less than 1100 we have the following:

$$J_1(4|p) = \{(p-1)/2, p-1\} \cup T(4|p) \cup \{p-1-r : r \in T(4|p)\}$$

where $T(4|p)$ are listed in Table 3 if $T(2|p) \neq \emptyset$. Moreover, $J_1(4|p^4) = J_1(4|p^2) = \emptyset$ for all primes in this range. As a consequence

$$J(4|p) = \{0, (p-1)/2, p-1\} \cup T(4|p) \cup \{p-1-r : r \in T(4|p)\} \quad \forall p < 1100.$$

Example 5. Let's turn to the partial sums of $\zeta(5)$ series. First we easily have $J_1(5|3) = \{2\}$ and $J_1(5|3^3) = \emptyset$ so that $J(5|3) = \{0, 2\}$. Furthermore

$$H(5;1) = 1,\ H(5;2) \equiv 44,\ H(5;3) \equiv 26,\ H(5;4) \equiv 1 \pmod{5^4}.$$

So we get $J_1(5|5) = J_1(5|5^2) = J_1(5|5^3) = \{4\}$, but $J_1(5|5^5) = \emptyset$. So $J(5|5) = \{0, 4\}$. Now we jump to $p = 37$ because it gives the first irregular pair $(37, 37-5)$ and therefore shows a different

p	$T(4\|p)$	p	$T(4\|p)$	p	$T(4\|p)$	p	$T(4\|p)$	p	$T(4\|p)$	p	$T(4\|p)$
17	2	157	61	307	32	487	125	757	115	941	341,355
41	18	163	10	311	89	499	97,137,232	773	354	947	241
59	15	181	74	317	11	503	60	787	356	967	90139
67	24	191	33	331	134	523	186	797	43	997	196
71	28	199	3,63	337	43	541	111	809	374	1009	383
79	6	227	43	359	53	617	51	827	320	1021	359
97	38	229	67	401	188	619	255	829	84	1031	141
101	46	239	42	409	99	647	40,194	839	88,238	1033	22
103	47	251	16,88	431	13	691	292	863	252	1039	107,391
131	58	257	112	461	209	701	160	883	212	1049	404
137	10,51	277	91	463	30	719	354	919	250	1061	212
139	63	283	11	479	54	743	281	937	89	1093	203

Table 3: The set $J(4|p) = \{0, (p-1)/2, p-1\} \cup T(4|p) \cup \{p-1-r : r \in T(4|p)\}$, $p \leq 1100$, is finite.

pattern. We must have $18 \in J_1(5|37)$ as guaranteed by Proposition 4.15(a). Computation shows that in fact $J_1(5|37) = \{6, 9, 12, 18, 24, 27, 30, 36\}$. We also have $J(5|37^2) = \{0, 6, 36\}$ which is a little strange, as the set is obviously asymmetric. Finally $J_1(5|37^5) = \emptyset$ which implies that

$$J(5|37) = \{0, 6, 9, 12, 18, 24, 27, 30, 36\}.$$

In general, when primes p ranges from 7 to 1061 but $p \neq 37$ we find by Maple

$$J(5|p) = \{0, p-1\} \cup T(5|p) \cup \{p-1-r : r \in T(5|p)\}, \quad J(5|p^2) = \{0, p-1\}, \quad J(5|p^3) = \emptyset,$$

where $T(5|p)$ is given by Table 4 whenever $T(5|p) \neq \emptyset$. The inclusion $p-1 \in J(5|p^2)$ is enforced by Lemma 2.2 and Theorem 2.6.

p	$T(5\|p)$	p	$T(5\|p)$	p	$T(5\|p)$	p	$T(5\|p)$	p	$T(5\|p)$	p	$T(5\|p)$
11	2	179	63	317	142,154	467	179	743	299,334	929	268
29	6	197	27	349	111	479	94,100	761	265	937	349
47	14	223	32,35	359	55	563	42	773	217	947	26
71	9	241	49	379	187	571	43	811	61	967	6,24
83	3,15,21	257	26	397	48	613	99	821	180	991	94
97	3,22	269	128	409	151	641	97	853	168	1009	66,451
107	37	283	38	419	95	659	137,286	857	274	1021	449
127	19	307	124	431	151	683	115	859	89	1031	139
149	14	311	144	433	197	691	273	877	324	1033	362
157	32	313	10,95	439	211	709	344	883	54	1039	181

Table 4: The set $J(5|p) = \{0, p-1\} \cup T(5|p) \cup \{p-1-r : r \in T(5|p)\}$, $p \leq 1061$, is finite.

Similar but larger range of data is available online [3]. To summarize we have

Proposition 6. *Let p be a prime such that $p \leq 3001$. Then $J(s|p)$ is finite for $2 \leq s \leq 300$.*

For other types of $\vec{s}$ we need the Criterion Theorem in our main article [2].

Criterion Theorem. *Let $d \geq 2$ be a positive integer and p be a prime such that $d \in G_{t_0}$. Let $\vec{s} = (s_1, \cdots, s_d) \in \mathbb{N}^d$ and put $m = \min\{s_i : 1 \leq i \leq d\}$. For $t \in \mathbb{N}$ set $f(\vec{s}, p; t) = \min\{-v_p\big(H(\vec{s}; n)\big) : n \in G_t\}$. If there is $\tau > t_0$ such that*

$$f(\vec{s}, p; \tau) > (\mathrm{wt}(\vec{s}) - m)(\tau - 1) - m,$$

then $J(\vec{s}|p)$ is finite.

Proposition 7. *Let s and d be two positive integers. Suppose $2 \leq s \leq 6$ and $d \leq 5$. Then the p-divisible set $J(\{s\}^d|p)$ is finite for the consecutive five primes immediately after $sd + 2$. Moreover there's always some prime p such that $J(\{s\}^d|p) = RJ(\{s\}^d; p)$ where*

$$RJ(\{s\}^d; p) = \begin{cases} \{0, p-1\} & \text{if } 2 \nmid s, \\ \{0, i + (p-1)/2, p-1, p : 0 \leq i \leq d-1\} & \text{if } 2 | s. \end{cases}$$

Proof. We may assume $d \geq 2$ by Prop. 6. Then the proposition follows from Criterion Theorem by computer computation whose results are listed in Tables 6-Ta:6dp. We put the Maple code at the end of this supplement. In each row of the tables we list the number τ used when applying Criterion Theorem and the discrepancy $J(\{s\}^d|p) \setminus RJ(\{s\}^d; p)$, separated by a semicolon. We include the data for $s = 1$ for future reference where we list RJ itself instead of the discrepancy when $p \leq d + 2$.

p	$d = 2$	$d = 3$	$d = 4$	$d = 5$
3	6, {0,5}	10, {0,8}	> 12, {0}	>10, {0,8}
7	4, {4}	>8, {11}	>8, $\emptyset$	> 8, {0,6,14}
11	>8 $\emptyset$	>8 $\emptyset$	>8, $\emptyset$	>8, $\emptyset$
13	4, $\emptyset$	>8, $\emptyset$	>8, $\emptyset$	>8, $\emptyset$
17	>8 {11,13}	>8, {5}	>8, $\emptyset$	>8, $\emptyset$
31	4, {17,22}			

Table 5: Finiteness of $J(\{1\}^d|p)$ for $2 \leq d \leq 5$.

p	$d = 2$	$d = 3$	$d = 4$	$d = 5$
7	3, $\emptyset$			
11	2, $\emptyset$	3, $\emptyset$	4, $\emptyset$	
13	2, {4}	3, $\emptyset$	4, $\emptyset$	5, $\emptyset$
17	2, $\emptyset$	3, {14}	4, $\emptyset$	5, $\emptyset$
19	2, $\emptyset$	3, {14}	4, $\emptyset$	5, $\emptyset$
23	2, {17}	3, $\emptyset$	4, $\emptyset$	5, $\emptyset$
29	2, $\emptyset$	3, $\emptyset$	4, {11}	5, $\emptyset$
31	2, $\emptyset$	4, {5,11}		
37	2, {6}			
41	2, $\emptyset$			

Table 6: Finiteness of $J(\{2\}^d|p)$ for $2 \leq d \leq 5$.

$\square$

p	$d=2$	$d=3$	$d=4$	$d=5$
11	2, $\emptyset$			
13	2, $\emptyset$	3, $\{10\}$		
17	2, $\{6\}$	3, $\{10,13\}$	4, $\emptyset$	
19	2, $\{12\}$	3, $\emptyset$	4, $\emptyset$	5, $\{10,16\}$
23	2, $\emptyset$	3, $\{7,15\}$	4, $\emptyset$	5, $\{14\}$
29	2, $\emptyset$	3, $\emptyset$	4, $\{6,9,23\}$	5, $\{11\}$
31	2, $\{11\}$		4, $\{18,28\}$	5, $\emptyset$
37	3, $\{13,14,21,23,24,28,73\}$			5, $\emptyset$

Table 7: Finiteness of $J(\{3\}^d|p)$ for $2 \leq d \leq 5$.

p	$d=2$	$d=3$	$d=4$	$d=5$
11	2, $\emptyset$			
13	2, $\{4,10\}$			
17	2, $\{13\}$	3, $\{6,13,14\}$		
19	2, $\emptyset$	3, $\emptyset$	4, $\emptyset$	
23	2, $\emptyset$	3, $\emptyset$	4, $\emptyset$	5, $\emptyset$
29	2, $\emptyset$	3, $\emptyset$	4, $\{24\}$	5, $\{11,24,25\}$
31	2, $\emptyset$	3, $\{21,24\}$	4, $\{28\}$	5, $\emptyset$
37	2, $\emptyset$		4, $\emptyset$	5, $\{31\}$
41	2, $\{40\}$			5, $\{8,18\}$

Table 8: Finiteness of $J(\{4\}^d|p)$, $2 \leq d \leq 5$.

p	$d=2$	$d=3$	$d=4$	$d=5$
13	2, $\emptyset$			
17	2, $\{7\}$			
19	2, $\{10\}$	3, $\{14\}$		
23	2, $\{3\}$	3, $\emptyset$	4, $\{10,15\}$	
29	2, $\emptyset$	3, $\emptyset$	4, $\{8,18\}$	5, $\{18,19\}$
31	2, $\emptyset$	3, $\{5,10,22,27\}$	4, $\{12,18,21\}$	5, $\emptyset$
37	2, $\{18,19\}$	3, $\{25,31\}$	4, $\{9,18\}$	5, $\{32\}$
41	2, $\{5,16\}$		4, $\{17,34,35\}$	5, $\{34,35,36\}$
43	2, $\{7,37\}$			5, $\emptyset$

Table 9: Finiteness of $J(\{5\}^d|p)$ for $2 \leq d \leq 5$.

p	$d=2$	$d=3$	$d=4$	$d=5$
17	2, $\emptyset$			
19	2, $\{4,7,16\}$			
23	2, $\emptyset$	3, $\{8\}$		
29	2, $\{24\}$	3, $\emptyset$	4, $\emptyset$	
31	2, $\{26\}$	3, $\{26,27\}$	4, $\{8,13,26\text{-}28\}$	
37	2, $\{16\}$	3, $\emptyset$	4, $\{9,15,23,33\}$	5, $\{9,10,15,16,33,34\}$
41	2, $\{28,35\}$	3, $\{18\}$	4, $\{32\}$	5, $\emptyset$
43	2, $\emptyset$		4, $\{10,12,37,38\}$	5, $\{10,11,14,18,29,37\text{-}39\}$
47	2, $\{34\}$			5, $\{36\}$
53	2, $\{37,49\}$			5, $\emptyset$

Table 10: Finiteness of $J(\{6\}^d|p)$, $2 \le d \le 5$.

Now we turn to non-homogeneous MZV series. We have the following results.

Proposition 8. *Let* $1 \le s \le 10$, $2 \le t \le 10$ *be two positive integers. Then there's always some prime* $p > sd + 2$ *such that* $J(s,t|p) = RJ(s,t;p)$ *where* $RJ(s,t)$ *is the reserved set for* $\zeta(s,t)$*.*

Proof. The proposition follows from Criterion Theorem by computer computation whose results are listed in Table 11. In each entry after (s,t) we list the prime p, the number τ used when applying Criterion Theorem to show finiteness of $J(s,t|p)$. For these primes we found that $J(s,t|p) = RJ(s,t;p)$.

s,t	$p;\tau$	s,t	$p;\tau$	s,t	$p;\tau$	s,t	$p;\tau$	s,t	$p;\tau$
1,1	13,4	2,1	13,4	3,1	11,>8	4,1	13,>8	5,1	13,>8
6,1	11,>6	7,1	19,>6	8,1	13,>6	9,1	17,>6	10,1	31,>6
1,2	11,4	2,2	7,3	3,2	11,3	4,2	11,3	5,2	19,4
6,2	13,4	7,2	19,5	8,2	13,5	9,2	23,6	10,2	31,6
1,3	7,5	2,3	41,3	3,3	11,2	4,3	11,3	5,3	11,3
6,3	13,3	7,3	13,4	8,3	13,4	9,3	13,4	10,3	17,4
1,4	11,4	2,4	11,3	3,4	17,3	4,4	11,2	5,4	19,3
6,4	13,3	7,4	17,3	8,4	17,3	9,4	19,3	10,4	17,3
1,5	11,4	2,5	11,3	3,5	11,3	4,5	17,3	5,5	13,2
6,5	47,3	7,5	19,3	8,5	17,3	9,5	17,3	10,5	19,3
1,6	13,4	2,6	11,3	3,6	17,3	4,6	13,3	5,6	43,3
6,6	17,2	7,6	17,3	8,6	31,3	9,6	19,3	10,6	19,3
1,7	17,4	2,7	29,3	3,7	19,3	4,7	37,3	5,7	17,3
6,7	19,3	7,7	17,2	8,7	29,3	9,7	23,3	10,7	23,3
1,8	17,4	2,8	13,3	3,8	19,3	4,8	23,3	5,8	17,3
6,8	17,3	7,8	41,3	8,8	19,2	9,8	29,3	10,8	23,3
1,9	19,4	2,9	17,3	3,9	17,3	4,9	17,3	5,9	17,3
6,9	19,3	7,9	29,3	8,9	31,3	9,9	29,2	10,9	29,3
1,10	19,4	2,10	19,3	3,10	19,3	4,10	31,3	5,10	29,3
6,10	29,3	7,10	23,3	8,10	23,3	9,10	23,3	10,10	23,2

Table 11: Finiteness of $J(s,t|p)$ for $2 \le s,t \le 5$.

$\square$

More data is available online [4].

Proposition 9. *Let $2 \leq r, s, t \leq 5$ be three positive integers. Then there's always some prime $p > sd + 2$ such that $J(r, s, t|p) = RJ(r, s, t; p)$ where $RJ(r, s, t)$ is the reserved set for $\zeta(r, s, t)$.*

Proof. The data of the proof is listed in Table 12. ☐

r,s,t	$p;\tau$	r,s,t	$p;\tau$	r,s,t	$p;\tau$	r,s,t	$p;\tau$	r,s,t	$p;\tau$
(1,1,1)	11,> 8	(1,1,2)	7,6	(1,1,3)	13,4	(1,1,4)	13,4	(1,1,5)	11,4
(1,2,1)	7,> 8	(1,2,2)	11,4	(1,2,3)	11,4	(1,2,4)	11,6	(1,2,5)	11,4
(1,3,1)	11,>6	(1,3,2)	11,>6	(1,3,3)	11,6	(1,3,4)	11,6	(1,3,5)	17,6
(1,4,1)	13,>6	(1,4,2)	19,>6	(1,4,3)	29,>6	(1,4,4)	13,>6	(1,4,5)	19,>6
(1,5,1)	11,>6	(1,5,2)	11,> 8	(1,5,3)	13,8	(1,5,4)	17,>6	(1,5,5)	37,>6
(2,1,1)	7,>9	(2,1,2)	47,>6	(2,1,3)	19,>6	(2,1,4)	11,>5	(2,1,5)	13,>6
(2,2,1)	17,6	(2,2,2)	11,3	(2,2,3)	11,3	(2,2,4)	13,3	(2,2,5)	13,3
(2,3,1)	23,>6	(2,3,2)	13,4	(2,3,3)	13,3	(2,3,4)	13,3	(2,3,5)	13,3
(2,4,1)	17,>6	(2,4,2)	17,4	(2,4,3)	13,4	(2,4,4)	29,5	(2,4,5)	19,4
(2,5,1)	17,>6	(2,5,2)	17,4	(2,5,3)	13,4	(2,5,4)	19,5	(2,5,5)	17,4
(3,1,1)	11,>6	(3,1,2)	11,>6	(3,1,3)	43,>6	(3,1,4)	17,>6	(3,1,5)	29,>6
(3,2,1)	11,>6	(3,2,2)	11,3	(3,2,3)	13,4	(3,2,4)	19,3	(3,2,5)	17,3
(3,3,1)	11,>6	(3,3,2)	13,5	(3,3,3)	19,4	(3,3,4)	23,3	(3,3,5)	17,3
(3,4,1)	11,>6	(3,4,2)	13,4	(3,4,3)	13,4	(3,4,4)	17,3	(3,4,5)	19,3
(3,5,1)	31,>6	(3,5,2)	17,6	(3,5,3)	17,4	(3,5,4)	17,4	(3,5,5)	29,4
(4,1,1)	11,>6	(4,1,2)	11,>6	(4,1,3)	17,>6	(4,1,4)	13,>6	(4,1,5)	13,>6
(4,2,1)	13,>6	(4,2,2)	23,5	(4,2,3)	13,4	(4,2,4)	37,4	(4,2,5)	37,5
(4,3,1)	31,>6	(4,3,2)	19,5	(4,3,3)	17,3	(4,3,4)	19,4	(4,3,5)	17,3
(4,4,1)	17,>6	(4,4,2)	23,6	(4,4,3)	17,4	(4,4,4)	19,3	(4,4,5)	17,3
(4,5,1)	13,>6	(4,5,2)	17,5	(4,5,3)	23,4	(4,5,4)	31,3	(4,5,5)	17,3
(5,1,1)	11,>6	(5,1,2)	13,>6	(5,1,3)	19,>6	(5,1,4)	29,>6	(5,1,5)	17,>6
(5,2,1)	11,>6	(5,2,2)	13,5	(5,2,3)	17,4	(5,2,4)	17,4	(5,2,5)	17,5
(5,3,1)	17,>6	(5,3,2)	17,6	(5,3,3)	17,4	(5,3,4)	17,4	(5,3,5)	17,4
(5,4,1)	13,>6	(5,4,2)	29,6	(5,4,3)	17,4	(5,4,4)	19,3	(5,4,5)	17,3
(5,5,1)	19,>6	(5,5,2)	17,6	(5,5,3)	47,4	(5,5,4)	23,3	(5,5,5)	23,3

Table 12: Finiteness of $J(r, s, t|p)$ for $2 \leq r, s, t \leq 5$.

More data is available online [4]. We have

Finally we make a conjecture about 2-divisible sets.

Conjecture 10. *For all positive integer d and $\vec{s} \in \mathbb{N}^d$ the 2-divisible set $J(\vec{s}|2) = \{0\}$.*

We put the supporting data in [5] due to its size. We also deal with prime 3 and 5 separately in [6, 7].

Proposition 11. *Let $p = 2, 3$ or 5. Then $J(\vec{s}|p)$ is finite for*

1. *$\vec{s} = (s, t)$ where $s, t \leq 20$ and $t \geq 2$.*
2. *$\vec{s} = (r, s, t)$ where $r, s, t \leq 10$ and $t \geq 2$.*
3. *$\vec{s} = (q, r, s, t)$ where $q, r, s, t \leq 4$ and $t \geq 2$.*

Appendix I: The program code

In this appendix we provide the pseudo code that we use to do the computation according to our main Criterion Theorem.

Let $\vec{s} = (s_1, \ldots, s_d)$, $m = \min\{s_i : 1 \leq i \leq d\}$, and p be a fixed prime. We want to use modular arithmetic because it's much faster than precise computation of the partial sums. So when

$n \in G_{t+1}$ we compute $p^{\mathrm{wt}(\vec{s})} H(\vec{s}; n)$. By the Criterion Theorem we see that if $p^{t\,\mathrm{wt}(\vec{s})} H(\vec{s}; n) \not\equiv 0 \pmod{p^{mt+m}}$ for all $n \in G_{t+1}$ then we can take $\tau = t+1$ and we're done.

Suppose we look for τ in the range of 2 to $e-1$ for some positive integer e. In the inner for-loop of the following pseudo code we use the formula

$$H(s_1, \dots, s_i; n-d+i) = H(s_1, \dots, s_i; n-1-d+i) + \frac{H(s_1, \dots, s_{i-1}; n-d+(i-1))}{(n-d+i)^{s_i}}.$$

```
Find_tau:=proc(s,p,e);

if p<d+1 then
  t:=floor(log(d)/log(p));
else
  t:=0;

m:=min(s[1],...,s[d]);
wt[i]:=s[1]+...+s[i] for i=1,...d;
psum[i]:=0 for i=1,...d;
ifstop=NO;

while t<e and ifstop=NO do
  ifstop:=YES;
  psum[i]:=psum[i]*p^wt[i] for i=1,...d;
  # In the beginning of each new segment G_{t+1}, the partial sums
  # H(...; p^t-1) should be multiplied by higher powers of p.

  for n from max(d,p^t) to p^(t+1)-1 do
    for i to d do
      psum[i]:=psum[i]+p^(t*s[i])*psum[i-1]/(n-d+i)^(s[i]) mod p^(m*t+m);
    end of inner for loop;
    if psum[d]=0 then ifstop:=NO;
  end of outer for loop;
  t:=t+1;
end while loop;

if ifstop=YES then
  tau:=t;
else
  e has to be larger.
```

Appendix II: Conjectural structure of the set $J_1(\vec{s}|p)$

The following conjecture is [1, Conj. 7.4].

Conjecture 12. *Suppose $\vec{s} \in \mathbb{N}^l$ such that (i) $\vec{s} = 1$, or (ii) $\vec{s} = (1,2,1)$, or (iii) $\vec{s} = (2r-1,1)$ for some $r \ge 1$, or (iv) $\vec{s} = 1^{2l}$ for some $l \ge 1$. Then $RJ(\vec{s}) = RJ_2(\vec{s})$. For all other $\vec{s}$ we have $RJ(\vec{s}) = RJ_1(\vec{s})$.*

Definition 13. Let $\vec{s} \in \mathbb{N}^l$. Define the *reserved density* of the MHS $H(\vec{s})$ by

$$\mathrm{density}(RJ(\vec{s}); X) = \frac{\sharp\{\text{prime } p : |\vec{s}| + 2 < p < X, J(\vec{s}|p) = RJ(\vec{s})\}}{\sharp\{\text{prime } p : |\vec{s}| + 2 < p < X\}} \tag{2}$$

and the mth *reserved density* by

$$\mathrm{density}(RJ^m(\vec{s}); X) = \frac{\sharp\{\text{prime } p : |\vec{s}| + 1 < p < X, \cup_{t=0}^{m} J_t(\vec{s}|p) = RJ_m(\vec{s})\}}{\sharp\{\text{prime } p : |\vec{s}| + 2 < p < X\}}. \tag{3}$$

The next conjecture is [1, Conj. 7.6].

Conjecture 14. *Let* $\vec{s} \in \mathbb{N}^l$. *Then*

$$\text{density}(RJ(\vec{s}); \infty) = \begin{cases} 1/\sqrt{e}, & \textit{if } l = 1, \vec{s} \geq 2, \\ 1/e, & \textit{if } l = \vec{s} = 1 \textit{ or } l \geq 2. \end{cases}$$

Assuming [1, Conj. 7.4] we have the following data to support [1, Conj. 7.6]. Throughout Tables 13-16 we set $d(\vec{s}) = \text{density}(RJ(\vec{s}); 3000)$.

References

[1] J. Zhao, Wolstenholme Type Theorem for multiple harmonic sums, *Intl. J. of Num. Thy.* **4**(1) (2008) 73-106.

[2] ______, *Finiteness of p-Divisible Sets of Multiple Harmonic Sums*, available online `http://arxiv.org/abs/math.NT/0303043v3`

[3] ______, *Finiteness Conjecture of* $J(s|p)$, `http://home.eckerd.edu/~zhaoj/research.htm`

[4] ______, *Finiteness Conjecture of* $J(s,t|p)$, $J(r,s,t|p)$ *and* $J(s,t,s,t|p)$, ibid.

[5] ______, *The Conjecture of* $J(\vec{s}|2) = \{0\}$, ibid.

[6] ______, *Finiteness of* $J(\vec{s}|3)$, ibid.

[7] ______, *Finiteness of* $J(\vec{s}|5)$, ibid.

$\vec{s}$	$d(\vec{s})$	$\vec{s}$	$d(\vec{s})$	$\vec{s}$	$d(\vec{s})$	$\vec{s}$	$d(\vec{s})$	$\vec{s}$	$d(\vec{s})$	$\vec{s}$	$d(\vec{s})$
2	63.40%	3	60.37%	4	55.84%	5	61.22%	6	64.87%	7	58.78%
8	58.55%	9	62.06%	10	60.33%	11	63.38%	12	54.12%	13	61.65%
14	60.47%	15	61.41%	16	62.97%	17	60.14%	18	61.47%	19	63.83%
20	62.65%	21	62.18%	22	60.19%	23	63.74%	24	59.24%	25	61.14%
26	59.24%	27	62.32%	28	60.10%	29	63.18%	30	63.10%	2,1	36.60%
3,1	35.51%	4,1	37.38%	5,1	38.88%	6,1	35.36%	7,1	34.19%	8,1	34.19%
9,1	34.74%	10,1	39.44%	3,2	39.25%	4,2	35.60%	5,2	38.17%	6,2	37.00%
7,2	40.05%	8,2	40.38%	9,2	40.38%	10,2	39.53%	4,3	35.36%	5,3	39.11%
6,3	36.77%	7,3	36.38%	8,3	38.73%	9,3	38.12%	10,3	39.29%	5,4	35.60%
6,4	39.91%	7,4	42.02%	8,4	37.41%	9,4	40.94%	10,4	40.00%	6,5	37.79%
7,5	38.35%	8,5	37.88%	9,5	34.35%	10,5	35.06%	7,6	33.41%	8,6	34.35%
9,6	41.18%	10,6	35.38%	8,7	36.24%	9,7	36.08%	10,7	37.26%	9,8	39.62%
10,8	42.08%	10,9	38.06%	1,2	34.96%	1,3	40.19%	2,3	36.22%	1,4	36.92%
2,4	36.77%	3,4	38.41%	1,5	39.34%	2,5	34.66%	3,5	40.28%	4,5	40.05%
1,6	38.41%	2,6	32.55%	3,6	39.58%	4,6	35.92%	5,6	37.56%	1,7	38.64%
2,7	35.60%	3,7	38.50%	4,7	37.09%	5,7	35.29%	6,7	36.94%	1,8	33.02%
2,8	34.74%	3,8	37.09%	4,8	38.82%	5,8	39.53%	6,8	37.41%	7,8	36.94%
1,9	35.45%	2,9	34.98%	3,9	39.53%	4,9	36.94%	5,9	35.53%	6,9	38.12%
7,9	34.67%	8,9	40.33%	1,10	34.27%	2,10	35.76%	3,10	38.12%	4,10	37.65%
5,10	35.29%	6,10	35.38%	7,10	40.09%	8,10	38.77%	9,10	33.33%	1,1	38.23%
2,2	39.72%	3,3	41.92%	4,4	33.26%	5,5	37.56%	6,6	37.65%	7,7	37.18%
8,8	36.08%	9,9	40.66%	10,10	37.35%	11,11	41.47%	12,12	35.31%	13,13	38.39%
14,14	44.18%	15,15	39.29%	16,16	36.90%	17,17	39.76%	18,18	35.08%	19,19	37.23%
20,20	38.04%	21,21	38.13%	22,22	34.53%	23,23	39.90%	24,24	35.82%	25,25	40.14%
26,26	40.24%	27,27	33.98%	28,28	37.11%	29,29	38.89%	30,30	35.35%	1,1,1	38.93%
1,2,1	39.02%	1,3,1	37.38%	1,4,1	37.47%	1,5,1	37.00%	1,6,1	34.43%	1,7,1	32.32%
1,8,1	40.84%	1,9,1	37.09%	1,10,1	36.94%	1,1,2	36.22%	1,2,2	36.92%	1,3,2	37.47%
1,4,2	37.24%	1,5,2	36.77%	1,6,2	34.19%	1,7,2	34.51%	1,8,2	36.15%	1,9,2	35.53%
1,10,2	31.06%	1,1,3	36.45%	1,2,3	37.70%	1,3,3	35.83%	1,4,3	39.11%	1,5,3	40.28%
1,6,3	36.38%	1,7,3	36.62%	1,8,3	36.47%	1,9,3	39.53%	1,10,3	39.76%	1,1,4	40.75%
1,2,4	36.77%	1,3,4	40.05%	1,4,4	38.88%	1,5,4	38.50%	1,6,4	37.32%	1,7,4	39.53%
1,8,4	37.88%	1,9,4	38.35%	1,10,4	38.59%	1,1,5	38.88%	1,2,5	37.47%	1,3,5	34.43%
1,4,5	40.14%	1,5,5	35.68%	1,6,5	37.65%	1,7,5	38.59%	1,8,5	34.59%	1,9,5	36.71%
1,10,5	35.61%	1,1,6	40.98%	1,2,6	37.70%	1,3,6	36.38%	1,4,6	35.68%	1,5,6	34.12%
1,6,6	39.76%	1,7,6	37.18%	1,8,6	36.71%	1,9,6	35.61%	1,10,6	40.09%	1,1,7	35.60%
1,2,7	38.97%	1,3,7	39.91%	1,4,7	36.94%	1,5,7	37.88%	1,6,7	38.59%	1,7,7	36.94%
1,8,7	37.50%	1,9,7	36.32%	1,10,7	40.19%	1,1,8	39.20%	1,2,8	39.67%	1,3,8	34.35%
1,4,8	33.88%	1,5,8	36.71%	1,6,8	37.41%	1,7,8	38.92%	1,8,8	37.50%	1,9,8	36.88%
1,10,8	38.77%	1,1,9	42.72%	1,2,9	36.00%	1,3,9	33.41%	1,4,9	37.18%	1,5,9	38.35%
1,6,9	33.73%	1,7,9	39.39%	1,8,9	35.93%	1,9,9	33.57%	1,10,9	39.48%	1,1,10	35.53%
1,2,10	36.94%	1,3,10	40.24%	1,4,10	38.12%	1,5,10	33.73%	1,6,10	37.74%	1,7,10	38.30%
1,8,10	33.57%	1,9,10	38.06%	1,10,10	34.28%	2,1,1	46.03%	2,2,1	37.85%	2,3,1	39.34%
2,4,1	38.64%	2,5,1	34.19%	2,6,1	35.60%	2,7,1	40.14%	2,8,1	43.19%	2,9,1	34.35%
2,10,1	40.00%	2,1,2	34.81%	2,2,2	42.86%	2,3,2	42.16%	2,4,2	38.88%	2,5,2	40.05%
2,6,2	37.32%	2,7,2	35.45%	2,8,2	39.06%	2,9,2	34.59%	2,10,2	33.88%	2,1,3	36.30%
2,2,3	38.41%	2,3,3	35.36%	2,4,3	34.43%	2,5,3	35.21%	2,6,3	40.14%	2,7,3	39.29%
2,8,3	34.12%	2,9,3	33.18%	2,10,3	34.12%	2,1,4	35.36%	2,2,4	36.77%	2,3,4	40.05%
2,4,4	35.45%	2,5,4	37.56%	2,6,4	37.18%	2,7,4	35.76%	2,8,4	38.35%	2,9,4	34.82%
2,10,4	38.44%	2,1,5	37.24%	2,2,5	35.60%	2,3,5	35.92%	2,4,5	37.79%	2,5,5	40.71%
2,6,5	35.53%	2,7,5	34.12%	2,8,5	34.12%	2,9,5	36.32%	2,10,5	37.26%	2,1,6	36.30%
2,2,6	38.50%	2,3,6	37.09%	2,4,6	38.12%	2,5,6	34.35%	2,6,6	36.00%	2,7,6	37.65%
2,8,6	33.73%	2,9,6	33.73%	2,10,6	36.41%	2,1,7	37.32%	2,2,7	37.79%	2,3,7	39.29%

Table 13: Density Conjecture.

$\vec{s}$	$d(\vec{s})$	$\vec{s}$	$d(\vec{s})$	$\vec{s}$	$d(\vec{s})$	$\vec{s}$	$d(\vec{s})$	$\vec{s}$	$d(\vec{s})$	$\vec{s}$	$d(\vec{s})$
2,4,7	31.06%	2,5,7	36.94%	2,6,7	36.47%	2,7,7	35.85%	2,8,7	33.02%	2,9,7	39.01%
2,10,7	36.64%	2,1,8	33.80%	2,2,8	36.24%	2,3,8	37.41%	2,4,8	36.47%	2,5,8	36.94%
2,6,8	40.80%	2,7,8	41.98%	2,8,8	32.39%	2,9,8	34.99%	2,10,8	34.99%	2,1,9	36.47%
2,2,9	41.65%	2,3,9	36.47%	2,4,9	38.12%	2,5,9	38.21%	2,6,9	35.38%	2,7,9	37.35%
2,8,9	37.12%	2,9,9	36.17%	2,10,9	34.99%	2,1,10	34.59%	2,2,10	36.94%	2,3,10	33.41%
2,4,10	39.39%	2,5,10	36.79%	2,6,10	38.53%	2,7,10	36.17%	2,8,10	36.88%	2,9,10	37.82%
2,10,10	35.07%	3,1,1	40.89%	3,2,1	40.98%	3,3,1	38.17%	3,4,1	42.39%	3,5,1	38.64%
3,6,1	34.27%	3,7,1	33.10%	3,8,1	35.76%	3,9,1	36.47%	3,10,1	38.12%	3,1,2	36.77%
3,2,2	34.90%	3,3,2	35.83%	3,4,2	35.83%	3,5,2	41.55%	3,6,2	34.51%	3,7,2	32.24%
3,8,2	38.35%	3,9,2	38.12%	3,10,2	35.76%	3,1,3	36.53%	3,2,3	35.83%	3,3,3	35.83%
3,4,3	39.20%	3,5,3	34.51%	3,6,3	37.18%	3,7,3	37.65%	3,8,3	34.59%	3,9,3	38.59%
3,10,3	34.67%	3,1,4	40.28%	3,2,4	41.22%	3,3,4	38.50%	3,4,4	35.45%	3,5,4	33.18%
3,6,4	35.76%	3,7,4	35.76%	3,8,4	36.47%	3,9,4	39.62%	3,10,4	41.27%	3,1,5	37.00%
3,2,5	37.09%	3,3,5	33.80%	3,4,5	36.47%	3,5,5	39.06%	3,6,5	40.47%	3,7,5	40.71%
3,8,5	38.92%	3,9,5	37.03%	3,10,5	37.12%	3,1,6	38.03%	3,2,6	38.26%	3,3,6	36.24%
3,4,6	40.71%	3,5,6	36.47%	3,6,6	35.29%	3,7,6	34.91%	3,8,6	40.80%	3,9,6	37.82%
3,10,6	36.41%	3,1,7	33.33%	3,2,7	37.65%	3,3,7	36.71%	3,4,7	38.82%	3,5,7	37.18%
3,6,7	41.98%	3,7,7	37.97%	3,8,7	34.99%	3,9,7	38.30%	3,10,7	38.06%	3,1,8	39.29%
3,2,8	37.18%	3,3,8	42.82%	3,4,8	34.82%	3,5,8	38.21%	3,6,8	38.21%	3,7,8	33.57%
3,8,8	34.99%	3,9,8	36.88%	3,10,8	34.28%	3,1,9	36.47%	3,2,9	34.59%	3,3,9	39.29%
3,4,9	33.73%	3,5,9	37.03%	3,6,9	36.41%	3,7,9	38.77%	3,8,9	38.30%	3,9,9	35.46%
3,10,9	35.07%	3,1,10	33.65%	3,2,10	35.76%	3,3,10	36.79%	3,4,10	36.08%	3,5,10	36.41%
3,6,10	40.19%	3,7,10	38.06%	3,8,10	35.93%	3,9,10	37.68%	3,10,10	38.39%	4,1,1	37.70%
4,2,1	40.52%	4,3,1	33.72%	4,4,1	38.41%	4,5,1	40.38%	4,6,1	34.51%	4,7,1	38.59%
4,8,1	36.47%	4,9,1	35.53%	4,10,1	36.94%	4,1,2	32.79%	4,2,2	36.77%	4,3,2	38.88%
4,4,2	34.98%	4,5,2	34.98%	4,6,2	34.12%	4,7,2	36.71%	4,8,2	36.24%	4,9,2	38.59%
4,10,2	41.04%	4,1,3	35.36%	4,2,3	38.17%	4,3,3	42.02%	4,4,3	40.14%	4,5,3	36.47%
4,6,3	36.00%	4,7,3	39.06%	4,8,3	37.41%	4,9,3	39.62%	4,10,3	39.86%	4,1,4	39.11%
4,2,4	34.74%	4,3,4	41.32%	4,4,4	35.29%	4,5,4	32.71%	4,6,4	37.88%	4,7,4	34.35%
4,8,4	37.26%	4,9,4	38.44%	4,10,4	37.82%	4,1,5	36.85%	4,2,5	36.15%	4,3,5	41.41*%
4,4,5	35.29%	4,5,5	34.35%	4,6,5	38.12%	4,7,5	36.56%	4,8,5	38.92%	4,9,5	38.77%
4,10,5	34.28%	4,1,6	34.04%	4,2,6	35.29%	4,3,6	38.35%	4,4,6	39.29%	4,5,6	37.41%
4,6,6	36.08%	4,7,6	34.43%	4,8,6	38.77%	4,9,6	34.52%	4,10,6	40.19%	4,1,7	36.24%
4,2,7	38.82%	4,3,7	37.88%	4,4,7	36.94%	4,5,7	40.57%	4,6,7	34.43%	4,7,7	38.06%
4,8,7	35.46%	4,9,7	37.59%	4,10,7	34.52%	4,1,8	36.71%	4,2,8	40.47%	4,3,8	37.88%
4,4,8	35.61%	4,5,8	39.86%	4,6,8	34.99%	4,7,8	34.28%	4,8,8	40.90%	4,9,8	32.62%
4,10,8	37.92%	4,1,9	36.47%	4,2,9	41.18%	4,3,9	35.85%	4,4,9	38.68%	4,5,9	34.28%
4,6,9	34.28%	4,7,9	36.41%	4,8,9	39.72%	4,9,9	36.26%	4,10,9	39.81%	4,1,10	37.88%
4,2,10	40.80%	4,3,10	32.08%	4,4,10	41.61%	4,5,10	38.06%	4,6,10	37.82%	4,7,10	36.17%
4,8,10	40.76%	4,9,10	39.57%	4,10,10	37.20%	5,1,1	36.77%	5,2,1	36.30%	5,3,1	37.94%
5,4,1	39.67%	5,5,1	38.97%	5,6,1	38.59%	5,7,1	38.35%	5,8,1	38.35%	5,9,1	39.29%
5,10,1	38.44%	5,1,2	37.70%	5,2,2	37.47%	5,3,2	36.38%	5,4,2	34.98%	5,5,2	38.35%
5,6,2	36.71%	5,7,2	33.65%	5,8,2	34.59%	5,9,2	35.61%	5,10,2	36.56%	5,1,3	37.24%
5,2,3	38.50%	5,3,3	36.15%	5,4,3	37.65%	5,5,3	34.35%	5,6,3	36.71%	5,7,3	37.65%
5,8,3	38.44%	5,9,3	34.91%	5,10,3	38.30%	5,1,4	38.97%	5,2,4	35.92%	5,3,4	38.35*%
5,4,4	41.41%	5,5,4	35.53%	5,6,4	37.18%	5,7,4	33.73%	5,8,4	37.50%	5,9,4	36.41%
5,10,4	40.19%	5,1,5	39.20%	5,2,5	36.94%	5,3,5	39.76%	5,4,5	39.76%	5,5,5	37.88%
5,6,5	34.91%	5,7,5	37.50%	5,8,5	39.01%	5,9,5	35.93%	5,10,5	34.75%	5,1,6	39.06%
5,2,6	39.76%	5,3,6	35.53%	5,4,6	42.12%	5,5,6	32.78%	5,6,6	38.92%	5,7,6	39.48%
5,8,6	37.12%	5,9,6	40.19%	5,10,6	38.30%	5,1,7	38.35%	5,2,7	35.53%	5,3,7	36.00%
5,4,7	38.21%	5,5,7	40.09%	5,6,7	35.93%	5,7,7	32.86%	5,8,7	37.82%	5,9,7	38.53%
5,10,7	38.39%	5,1,8	36.24%	5,2,8	37.65%	5,3,8	34.67%	5,4,8	38.21%	5,5,8	40.19%

Table 14: Density Conjecture.

$\vec{s}$	$d(\vec{s})$	$\vec{s}$	$d(\vec{s})$	$\vec{s}$	$d(\vec{s})$	$\vec{s}$	$d(\vec{s})$	$\vec{s}$	$d(\vec{s})$	$\vec{s}$	$d(\vec{s})$
5,6,8	38.06%	5,7,8	37.82%	5,8,8	35.93%	5,9,8	41.00%	5,10,8	34.60%	5,1,9	36.24%
5,2,9	35.85%	5,3,9	34.67%	5,4,9	36.88%	5,5,9	37.12%	5,6,9	37.35%	5,7,9	39.95%
5,8,9	39.10%	5,9,9	37.44%	6,1,1	35.60%	6,2,1	36.30%	6,3,1	39.91%	6,4,1	37.56%
6,5,1	36.00%	6,6,1	36.94%	6,7,1	35.29%	6,8,1	39.06%	6,10,1	32.08%	6,9,1	37.50%
6,10,3	35.93%	6,1,2	35.83%	6,2,2	38.73%	6,3,2	38.73%	6,4,2	36.00%	6,5,2	37.41%
6,6,2	34.59%	6,7,2	39.29%	6,8,2	36.79%	6,9,2	37.74%	6,10,2	38.53%	6,1,3	34.98%
6,2,3	38.73%	6,3,3	35.76%	6,4,3	42.35%	6,5,3	38.12%	6,6,3	37.41%	6,7,3	32.78%
6,8,3	39.15%	6,9,3	34.99%	6,10,3	35.93%	6,1,4	38.03%	6,2,4	34.59%	6,3,4	34.35%
6,4,4	36.94%	6,5,4	40.71%	6,6,4	35.38%	6,7,4	37.03%	6,8,4	36.17%	6,9,4	37.35%
6,10,4	38.77%	6,1,5	35.53%	6,2,5	35.53%	6,3,5	35.06%	6,4,5	39.29%	6,5,5	41.51%
6,6,5	37.50%	6,7,5	33.33%	6,8,5	37.59%	6,9,5	32.62%	6,10,5	35.46%	6,1,6	39.53%
6,2,6	35.53%	6,3,6	40.00%	6,4,6	38.68%	6,5,6	32.55%	6,6,6	33.81%	6,7,6	40.43%
6,8,6	34.75%	6,9,6	37.35%	6,10,6	37.44%	6,1,7	37.18%	6,2,7	38.12%	6,3,7	36.08%
6,4,7	38.68%	6,5,7	39.01%	6,6,7	36.41%	6,7,7	30.50%	6,8,7	38.06%	6,9,7	36.73%
6,10,7	37.92%	6,1,8	35.53%	6,2,8	38.68%	6,3,8	39.39%	6,4,8	36.41%	6,5,8	39.48%
6,6,8	37.59%	6,7,8	38.77%	6,8,8	37.68%	6,9,8	40.76%	6,10,8	39.81%	6,1,9	38.21%
6,2,9	37.50%	6,3,9	37.82%	6,4,9	39.24%	6,5,9	39.48%	6,6,9	35.46%	6,7,9	35.31%
6,8,9	37.92%	6,9,9	33.41%	6,10,9	37.20%	6,1,10	38.68%	6,2,10	35.22%	6,3,10	39.01%
6,4,10	39.95%	6,5,10	35.70%	6,6,10	36.49%	6,7,10	35.54%	6,8,10	35.07%	6,9,10	36.97%
6,10,10	40.05%	7,1,1	42.62%	7,2,1	39.67%	7,3,1	36.38%	7,4,1	38.12%	7,5,1	39.29%
7,6,1	38.59%	7,7,1	35.76%	7,8,1	37.97%	7,9,1	32.08%	7,10,1	36.41%	7,1,2	38.97%
7,2,2	32.16%	7,3,2	40.94%	7,4,2	35.76%	7,5,2	33.88%	7,6,2	34.82%	7,7,2	35.14%
7,8,2	39.86%	7,9,2	35.22%	7,10,2	34.28%	7,1,3	38.03%	7,2,3	35.29%	7,3,3	36.47%
7,4,3	35.29%	7,5,3	34.82%	7,6,3	38.92%	7,7,3	36.32%	7,8,3	39.24%	7,9,3	35.93%
7,10,3	40.43%	7,1,4	39.53%	7,2,4	38.82%	7,3,4	38.35%	7,4,4	37.65%	7,5,4	35.61%
7,6,4	37.03%	7,7,4	37.59%	7,8,4	42.79%	7,9,4	36.17%	7,10,4	40.90%	7,1,5	34.59%
7,2,5	33.41%	7,3,5	35.76%	7,4,5	36.32%	7,5,5	41.27%	7,6,5	38.30%	7,7,5	36.41%
7,8,5	36.88%	7,9,5	40.90%	7,10,5	43.60%	7,1,6	35.76%	7,2,6	37.88%	7,3,6	43.40%
7,4,6	36.08%	7,5,6	35.70%	7,6,6	33.81%	7,7,6	39.01%	7,8,6	36.64%	7,9,6	33.89%
7,10,6	35.07%	7,1,7	34.12%	7,2,7	33.73%	7,3,7	39.39%	7,4,7	40.66%	7,5,7	36.64%
7,6,7	39.48%	7,7,7	37.59%	7,8,7	41.00%	7,9,7	39.10%	7,10,7	41.23%	7,1,8	37.03%
7,2,8	35.85%	7,3,8	35.70%	7,4,8	34.75%	7,5,8	38.06%	7,6,8	33.33%	7,7,8	37.20%
7,8,8	37.92%	7,9,8	40.52%	7,10,8	40.28%	7,1,9	38.68%	7,2,9	33.57%	7,3,9	37.59%
7,4,9	34.04%	7,5,9	31.44%	7,6,9	36.97%	7,7,9	33.18%	7,8,9	33.18%	7,9,9	33.18%
7,10,9	37.68%	7,1,10	35.22%	7,2,10	37.59%	7,3,10	36.64%	7,4,10	34.75%	7,5,10	34.36%
7,6,10	36.26%	7,7,10	35.54%	7,8,10	38.39%	7,9,10	35.54%	7,10,10	37.92%	8,1,1	35.92%
8,2,1	38.73%	8,3,1	36.94%	8,4,1	37.65%	8,5,1	36.24%	8,6,1	34.82%	8,7,1	35.14%
8,8,1	37.74%	8,9,1	33.81%	8,10,1	36.88%	8,1,2	36.62%	8,2,2	35.53%	8,3,2	41.18%
8,4,2	36.24%	8,5,2	39.29%	8,6,2	37.50%	8,7,2	37.50%	8,8,2	35.93%	8,9,2	35.70%
8,10,2	39.95%	8,1,3	41.65%	8,2,3	36.47%	8,3,3	39.76%	8,4,3	37.18%	8,5,3	37.74%
8,6,3	37.74%	8,7,3	37.12%	8,8,3	34.99%	8,9,3	39.95%	8,10,3	37.82%	8,1,4	35.06%
8,2,4	36.94%	8,3,4	39.29%	8,4,4	37.03%	8,5,4	37.03%	8,6,4	38.77%	8,7,4	40.19%
8,8,4	36.88%	8,9,4	34.28%	8,10,4	42.89%	8,1,5	36.47%	8,2,5	41.18%	8,3,5	36.79%
8,4,5	37.97%	8,5,5	36.17%	8,6,5	41.84%	8,7,5	36.17%	8,8,5	38.53%	8,9,5	38.15%
8,10,5	36.26%	8,1,6	35.76%	8,2,6	34.43%	8,3,6	35.38%	8,4,6	32.62%	8,5,6	41.37%
8,6,6	33.33%	8,7,6	36.17%	8,8,6	36.97%	8,9,6	38.15%	8,10,6	37.68%	8,1,7	33.73%
8,2,7	38.21%	8,3,7	35.70%	8,4,7	33.57%	8,5,7	36.64%	8,6,7	37.59%	8,7,7	31.04%
8,8,7	35.54%	8,9,7	34.83%	8,10,7	36.49%	8,1,8	39.86%	8,2,8	37.35%	8,3,8	39.01%
8,4,8	33.57%	8,5,8	37.35%	8,6,8	33.41%	8,7,8	36.97%	8,8,8	39.81%	8,9,8	36.02%
8,10,8	40.05%	8,1,9	39.72%	8,2,9	35.46%	8,3,9	35.93%	8,4,9	40.19%	8,5,9	34.83%
8,6,9	34.60%	8,7,9	35.78%	8,8,9	36.73%	8,9,9	39.57%	8,10,9	37.92%	8,1,10	34.04%
8,2,10	39.48%	8,3,10	34.04%	8,4,10	36.97%	8,5,10	33.65%	8,6,10	35.54%	8,7,10	36.49%

Table 15: Density Conjecture.

$\vec{s}$	$d(\vec{s})$	$\vec{s}$	$d(\vec{s})$	$\vec{s}$	$d(\vec{s})$	$\vec{s}$	$d(\vec{s})$	$\vec{s}$	$d(\vec{s})$	$\vec{s}$	$d(\vec{s})$
8,8,10	36.97%	8,9,10	38.63%	8,10,10	38.48%	9,1,1	40.38%	9,2,1	36.24%	9,3,1	35.76%
9,4,1	40.71%	9,5,1	36.71%	9,6,1	35.85%	9,7,1	33.49%	9,8,1	35.46%	9,9,1	34.28%
9,10,1	39.95%	9,1,2	37.88%	9,2,2	37.65%	9,3,2	32.24%	9,4,2	37.41%	9,5,2	40.09%
9,6,2	36.79%	9,7,2	35.70%	9,8,2	39.01%	9,9,2	39.72%	9,10,2	38.77%	9,1,3	40.47%
9,2,3	41.41%	9,3,3	37.18%	9,4,3	37.50%	9,5,3	36.08%	9,6,3	37.35%	9,7,3	34.52%
9,8,3	37.12%	9,9,3	37.59%	9,10,3	34.12%	9,1,4	34.82%	9,2,4	36.24%	9,3,4	37.50%
9,4,4	34.43%	9,5,4	34.99%	9,6,4	37.35%	9,7,4	36.88%	9,8,4	39.01%	9,9,4	34.12%
9,10,4	37.20%	9,1,5	35.29%	9,2,5	40.80%	9,3,5	37.26%	9,4,5	35.46%	9,5,5	39.01%
9,6,5	40.66%	9,7,5	35.46%	9,8,5	36.49%	9,9,5	37.68%	9,10,5	39.81%	9,1,6	34.20%
9,2,6	40.09%	9,3,6	34.75%	9,4,6	38.30%	9,5,6	35.22%	9,6,6	33.81%	9,7,6	38.15%
9,8,6	37.20%	9,9,6	36.97%	9,10,6	39.10%	9,1,7	38.21%	9,2,7	37.35%	9,3,7	33.10%
9,4,7	35.70%	9,5,7	39.24%	9,6,7	35.78%	9,7,7	38.39%	9,8,7	37.20%	9,9,7	42.42%
9,10,7	38.39%	9,1,8	35.93%	9,2,8	40.43%	9,3,8	35.46%	9,4,8	36.17%	9,5,8	36.97%
9,6,8	38.63%	9,7,8	37.68%	9,8,8	39.57%	9,9,8	36.26%	9,10,8	33.65%	9,1,9	33.57%
9,2,9	35.70%	9,3,9	39.24%	9,4,9	38.86%	9,5,9	40.28%	9,6,9	34.83%	9,7,9	34.60%
9,8,9	35.31%	9,9,9	36.97%	9,10,9	39.19%	9,1,10	33.33%	9,2,10	37.82%	9,3,10	36.97%
9,4,10	40.76%	9,5,10	38.63%	9,6,10	36.73%	9,7,10	36.02%	9,8,10	38.15%	9,9,10	37.29%
9,10,10	36.34%	10,1,1	36.71%	10,2,1	34.12%	10,3,1	37.18%	10,4,1	32.47%	10,5,1	40.33%
10,6,1	40.57%	10,7,1	36.64%	10,8,1	30.02%	10,9,1	39.95%	10,10,1	36.64%	10,1,2	36.47%
10,2,2	36.71%	10,3,2	36.71%	10,4,2	42.22%	10,5,2	41.04%	10,6,2	38.30%	10,7,2	35.22%
10,8,2	37.12%	10,9,2	38.30%	10,10,2	41.47%	10,1,3	36.47%	10,2,3	36.94%	10,3,3	39.15%
10,4,3	37.26%	10,5,3	37.82%	10,6,3	40.90%	10,7,3	36.88%	10,8,3	35.46%	10,9,3	38.86%
10,10,3	36.97%	10,1,4	34.82%	10,2,4	34.91%	10,3,4	39.39%	10,4,4	39.01%	10,5,4	35.22%
10,6,4	36.17%	10,7,4	35.93%	10,8,4	37.92%	10,9,4	33.18%	10,10,4	38.39%	10,1,5	35.14%
10,2,5	37.26%	10,3,5	34.99%	10,4,5	37.12%	10,5,5	35.70%	10,6,5	37.35%	10,7,5	35.54%
10,8,5	34.36%	10,9,5	38.15%	10,10,5	40.76%	10,1,6	36.32%	10,2,6	42.32%	10,3,6	34.52%
10,4,6	34.75%	10,5,6	36.17%	10,6,6	37.68%	10,7,6	39.34%	10,8,6	33.18%	10,9,6	36.73%
10,10,6	35.78%	10,1,7	38.77%	10,2,7	37.12%	10,3,7	38.30%	10,4,7	39.24%	10,5,7	36.73%
10,6,7	36.97%	10,7,7	34.60%	10,8,7	33.41%	10,9,7	35.54%	10,10,7	40.76%	10,1,8	31.68%
10,2,8	35.46%	10,3,8	35.22%	10,4,8	32.23%	10,5,8	38.15%	10,6,8	36.26%	10,7,8	37.44%
10,8,8	36.26%	10,9,8	35.78%	10,10,8	34.20%	10,1,9	39.24%	10,2,9	36.17%	10,3,9	32.46%
10,4,9	37.44%	10,5,9	39.57%	10,6,9	37.44%	10,7,9	34.83%	10,8,9	33.89%	10,9,9	34.92%
10,10,9	32.07%	10,1,10	37.12%	10,2,10	41.23%	10,3,10	36.26%	10,4,10	42.18%	10,5,10	35.54%
10,6,10	39.34%	10,7,10	34.36%	10,8,10	36.34%	10,9,10	38.24%	10,10,10	40.00%	$\{2\}^4$	35.60%
$\{4\}^4$	35.85%	$\{6\}^4$	33.89%	$\{8\}^4$	36.67%	$\{10\}^4$	37.08%	$\{12\}^4$	32.45%	$\{14\}^4$	41.20%
$\{16\}^4$	34.14%	$\{18\}^4$	38.05%	$\{20\}^4$	36.92%	$\{2\}^5$	37.79%	$\{4\}^5$	39.72%	$\{6\}^5$	35.71%
$\{8\}^5$	39.95%	$\{10\}^5$	41.11%	$\{12\}^5$	37.29%	$\{14\}^5$	33.82%	$\{16\}^5$	35.70%	$\{18\}^5$	37.10%
$\{20\}^5$	35.56%	$\{2\}^6$	39.76%	$\{4\}^6$	39.34%	$\{6\}^6$	32.70%	$\{8\}^6$	40.14%	$\{10\}^6$	38.02%
$\{12\}^6$	39.27%	$\{14\}^6$	40.93%	$\{16\}^6$	39.90%	$\{18\}^6$	36.32%	$\{20\}^6$	36.16%	$\{2\}^7$	36.00%
$\{4\}^7$	39.43%	$\{6\}^7$	36.21%	$\{8\}^7$	35.90%	$\{10\}^7$	36.98%	$\{2\}^8$	42.69%	$\{4\}^8$	35.00%
$\{6\}^8$	33.89%	$\{8\}^8$	39.71%	$\{10\}^8$	39.36%	$\{12\}^8$	35.22%	$\{14\}^8$	35.91%	$\{16\}^8$	36.25%
$\{18\}^8$	32.75%	$\{20\}^8$	36.80%	$\{2\}^9$	42.79%	$\{4\}^9$	35.08%	$\{6\}^9$	33.98%	$\{8\}^9$	39.76%
$\{10\}^9$	39.31%	$\{12\}^9$	35.57%	$\{14\}^9$	36.00%	$\{16\}^9$	36.52%	$\{18\}^9$	32.57%	$\{20\}^9$	36.50%
$\{2\}^{10}$	42.79%	$\{4\}^{10}$	35.17%	$\{6\}^{10}$	34.14%	$\{8\}^{10}$	39.61%	$\{10\}^{10}$	39.51%	$\{12\}^{10}$	35.66%
$\{14\}^{10}$	36.27%	$\{16\}^{10}$	36.29%	$\{18\}^{10}$	32.65%	$\{20\}^{10}$	36.10%				

Table 16: Density Conjecture.